

Eye Associates of South Georgia  
(229)247-4114

**PRK POST-OP INSTRUCTIONS:**

**GATIFLOXACIN:** Instill one drop **FOUR** times a day to: **RIGHT EYE/LEFT EYE/BOTH EYES**

**PREDNISOLONE:** Instill one drop **FOUR** times a day to:  
**RIGHT EYE/LEFT EYE/BOTH EYES**

- Instill **Gatifloxacin** drop in appropriate eye(s). Gently close eye(s) for 15 seconds. After 5 minutes, use **Prednisolone** drop in appropriate eye(s). At each post-op visit, the doctor will inform you on how long to continue using each eye drop.
- Goggles **MUST** be worn **ALL DAY TODAY AND SLEEP WITH TONIGHT**. In the morning you can remove goggles, but goggles **MUST** be worn for 1 week while sleeping.
- Use Systane preservative-free or a preservative-free artificial tear as often as needed, but **MUST** be preservative-free.
- Use **ketorolac** eye drops **three** times a day in operated eye.
- Take **Vitamin C** capsules 1,000 mg **once** a day for 60 days.
- Take 1-2 **Norco** tablets every 4-6 hours or as needed for pain.
- If you experience any pain or decreased vision, call the office phone number listed above.